WhatsApp

Botox / Filler

Post-Procedure Precautions



General Progress & Precautions

- •The effects of Botox can sometimes be felt the very next day, but generally appear around 3 days after treatment and peak at about 2–3 weeks. After that, the effects gradually diminish, and by about 1 month, slight expression lines may return, giving a more natural look.
- •Localized bruising may occur after Botox treatment, but this typically resolves within 7–10 days and is not a cause for concern.
- •If Botox was applied around the eyes, avoid lying down for about 4 hours after the procedure to help prevent eyelid drooping.

Wrinkle Botox

- •After treatment around the eyes, your smile may feel slightly unnatural at first.
- •Botox on the forehead may cause a heavy feeling in the eyebrows, slight discomfort when opening your eyes, or a drooping effect on the eyebrows. These effects usually improve as the treatment settles.

Jaw (Masseter) Botox

- •Botox works by reducing muscle activity. For jaw Botox, effects appear after about 3–4 weeks and reach their peak after 2–3 months.
- •If a second treatment is needed, it is usually scheduled 4–6 months after the first. Please visit on your scheduled date to assess your condition and receive additional treatment if necessary.
- •After square jaw Botox, avoiding activities that overuse the jaw muscles—such as chewing gum, eating tough foods like squid or jerky—will help maintain the smaller muscle size for longer.
- •In some cases, you may feel weaker chewing power for about 1 month after the procedure, but this typically returns to normal with time.

Filler

- •Bruising may occur after filler treatment, but it usually resolves within 7–10 days and is not a cause for concern.
- •Take any prescribed medications regularly and at the scheduled times.
- •Swelling is common after filler injections, but within about 2 weeks the treated area will settle into the desired contour.
- •If you experience symptoms such as heat, severe swelling, persistent pain, changes in skin color, blistering, or pustules, please contact the clinic and visit immediately.

X Please note that individual recovery may vary, and the guidance provided by your medical team may differ from the instructions above.



T. +82 10 3141 4711

A. B1, Aratower, 3, Seocho-daero 77-gil, Seocho-gu, Seoul W. made-young.co